

PARTITIONS WITH DESIGNATED SUMMANDS

G.E. ANDREWS, R.P. LEWIS, J. LOVEJOY

ABSTRACT. This paper is devoted to the study of a new class of partitions, partitions with designated summands. In such partitions, among those parts of the same magnitude one is tagged or designated. The resulting partition functions are related to several classical ordinary partition functions. Also they have many interesting divisibility properties and satisfy quite appealing modular relations.

1. INTRODUCTION

The object of our study is partitions with designated summands. They are constructed by taking ordinary partitions and tagging exactly one of each part size. Thus there are 15 partitions of 5 with designated summands:

$$5', 4' + 1', 3' + 2', 3' + 1' + 1, 3' + 1 + 1', 2' + 2 + 1', \\ 2 + 2' + 1', 2' + 1' + 1 + 1, 2' + 1 + 1' + 1, 2' + 1 + 1 + 1', 1' + 1 + 1 + 1 + 1, \\ 1 + 1' + 1 + 1 + 1, 1 + 1 + 1' + 1 + 1, 1 + 1 + 1 + 1' + 1, 1 + 1 + 1 + 1 + 1'.$$

The total number of partitions of n with designated summands is denoted by $PD(n)$. Hence $PD(5) = 15$.

We shall also study $PDO(n)$ the total number of partitions of n with designated summands in which all parts are odd. Thus $PDO(5) = 8$.

While these objects have not appeared in the literature before, they are tacitly considered by P. A. MacMahon [13] in his work on generalized divisor sums. Indeed, MacMahon's $A_{n,k}$ is the number of partitions of n with designated summands wherein exactly k different magnitudes occur among all the parts. MacMahon [13, Section 17] is able to connect $A_{n,k}$ with numerous divisor sum identities due to Glaisher [7], Ramanujan [15] and others. However, MacMahon neglected the very interesting case wherein the number of different magnitudes of parts is suppressed. We shall consider the generating functions:

$$(1.1) \quad pd(q) = \sum_{n=0}^{\infty} PD(n)q^n,$$

Date: January 21, 2004.

1991 Mathematics Subject Classification. 05A17, 11P81.

and

$$(1.2) \quad pdo(q) = \sum_{n=0}^{\infty} PDO(n)q^n.$$

In Section 2, we shall show (among other things) that

$$(1.3) \quad pd(q) = \frac{\eta(6z)}{\eta(z)\eta(2z)\eta(3z)},$$

where

$$(1.4) \quad \eta(z) = q^{\frac{1}{24}} \prod_{n=1}^{\infty} (1 - q^n),$$

and

$$(1.5) \quad q := e^{2\pi iz},$$

Also

$$(1.6) \quad pdo(q) = \frac{\eta(4z)\eta^2(6z)}{\eta(z)\eta(3z)\eta(12z)}.$$

We examine q -series identities in Section 3, and in Section 4 we relate $PD(n)$ and $PDO(n)$ to several well-known partition functions. As a result, we can easily deduce that $PDO(n)$ is even unless either n is 0 or n is a perfect square not divisible by 3.

Section 5 is devoted to arithmetic properties of partitions with designated summands and Section 6 we investigate some modular relations.

2. THE GENERATING FUNCTIONS

Theorem 1. *Suppose S is a set of positive integers. We denote by $PD_S(n)$ the number of partitions of n with designated summands all of whose parts lie in S , and we denote by $pd_S(q)$ the generating function for $PD_S(n)$. Then*

$$(2.1) \quad pd_S(q) = \prod_{n \in S} \frac{(1 - q^{6n})}{(1 - q^n)(1 - q^{2n})(1 - q^{3n})}.$$

Proof: We see immediately that

$$\begin{aligned}
pd_S(q) &= \prod_{n \in S} (1 + q^n + 2q^{2n} + 3q^{3n} + 4q^{4n} + \cdots) \\
&= \prod_{n \in S} \left(1 + \frac{q^n}{(1 - q^n)^2} \right) \\
&= \prod_{n \in S}^{\infty} \frac{(1 - q^n + q^{2n})}{(1 - q^n)^2} \\
&= \prod_{n \in S}^{\infty} \frac{(1 + q^{3n})}{(1 - q^n)^2(1 + q^n)} \\
&= \prod_{n \in S}^{\infty} \frac{(1 - q^{6n})}{(1 - q^n)(1 - q^{2n})(1 - q^{3n})}.
\end{aligned}$$

□

Corollary 2. Equation (1.3) is valid.

Proof: This is the case $S = \mathbb{N}$, the set of all positive integers. □

Corollary 3. If S_k is the set of all positive integers not divisible by k , then

$$pd_{S_k}(q) = pd(q)/pd(q^k).$$

Proof: This assertion is the specialization of Theorem 1 to S_k . □

Corollary 4. Equation (1.6) is valid.

Proof: This follows directly from Corollary 3 with $k = 2$. □

Indeed we might note that if S consists of all positive integers congruent to one of

a_1, a_2, \dots, a_j modulo k , and if S is symmetric (i.e. whenever $\alpha \in S$ then all positive integers $\equiv -\alpha \pmod{k}$ are in S), then $pd_S(q)$ is a modular function. The subsequent sections are restricted to the instances of $pd_S(q)$ that seem to have the most interest, namely $pd(q)$ and $pdo(q)$.

3. q -SERIES FOR $pd(q)$ AND $pdo(q)$

There are several reasons for considering q -series expansions of these generating functions. Often such expansions provide efficient algorithms for the calculation of their coefficients. This turns out to be the case for both $pd(q)$ (see (3.4)) and $pdo(q)$ (see (3.13)). In addition, we occasionally uncover new mysteries. In our present study, it is quite mysterious that the q -series (3.3) and (3.12) which seem to have complex coefficients, in fact do not.

Theorem 5. For $|q| < 1$ and $\zeta = e^{\pi i/3}$,

$$(3.1) \quad pd(q) = \frac{(\zeta q; q)_\infty (\zeta^{-1} q; q)_\infty}{(q; q)_\infty^2}$$

$$(3.2) \quad = 1 + \sum_{n=1}^{\infty} \frac{(-q^3; q^3)_{n-1} q^n}{(q; q)_n (q^2; q^2)_{n-1} (1 - q^n)}$$

$$(3.3) \quad = 1 + \zeta \sum_{n=1}^{\infty} \left(\frac{(\zeta q; q)_{n-1}}{(q; q)_n} \right)^3 (1 - \zeta q^{2n}) (-1)^n q^{\binom{n+1}{2}} \zeta^{-n}$$

$$(3.4) \quad = \frac{\sum_{n=0}^{\infty} q^{\binom{n+1}{2}} - 3 \sum_{n=0}^{\infty} q^{\binom{3n+2}{2}}}{\sum_{n=0}^{\infty} (-1)^n (2n+1) q^{\binom{n+1}{2}}},$$

where

$$(3.5) \quad (a; q)_n = (1 - a)(1 - aq) \cdots (1 - aq^{n-1}).$$

Proof: By (1.3),

$$\begin{aligned} pd(q) &= \frac{(q^6; q^6)_\infty}{(q; q)_\infty (q^2; q^2)_\infty (q^3; q^3)_\infty} \\ &= \frac{(-q^3; q^3)_\infty}{(q; q)_\infty^2 (-q; q)_\infty} \\ &= \frac{(\zeta q; q)_\infty (\zeta^{-1} q; q)_\infty}{(q; q)_\infty^2}, \end{aligned}$$

which proves (3.1). Identity (3.2) may be rewritten as follows:

$$(3.6) \quad \sum_{n=0}^{\infty} \frac{(\zeta; q)_n (\zeta^{-1}; q)_n}{(q; q)_\infty (q; q)_n} = \frac{(\zeta q; q)_\infty (\zeta^{-1} q; q)_\infty}{(q; q)_\infty (q; q)_\infty},$$

and in this form we see that it is a specialization of the q -analog of Gauss's theorem [2, p. 20, Cor. 2.4, $a = \zeta$, $b = \zeta^{-1}$, $c = q$]. Identity (3.3) may be rewritten as follows:

$$(3.7) \quad \lim_{t \rightarrow 0} \sum_{n=0}^{\infty} \frac{(\zeta; q)_n^3 (1 - \zeta q^{2n}) (t^{-1}; q)_n}{(q; q)_n^3 (1 - \zeta) (\zeta q t; q)_n} \left(\frac{t \zeta q}{\zeta^2} \right)^n = \frac{(\zeta^{-1} q; q)_\infty (\zeta q; q)_\infty}{(q; q)_\infty (q; q)_\infty},$$

and in this form we see that it is a specialization of the limiting form of Jackson's theorem [6, p. 238, eq. (II,20)].

To obtain (3.4), we first note that

$$(3.8) \quad \frac{(-1)^n(\zeta^{-n} - \zeta^{n+1})}{(1 - \zeta)} = \begin{cases} 1 & \text{if } n \equiv 0 \pmod{3} \\ -2 & \text{if } n \equiv 1 \pmod{3} \\ 1 & \text{if } n \equiv 2 \pmod{3}. \end{cases}$$

Hence by Jacobi's triple product [6, p. 239, eq. (II.28)]

$$\begin{aligned} \frac{(\zeta q; q)_\infty (\zeta^{-1} q; q)_\infty}{(q; q)_\infty^2} &= \frac{(q; q)_\infty (\zeta; q)_\infty (\zeta^{-1} q; q)_\infty}{(1 - \zeta)(q; q)_\infty^3} \\ &= \frac{\sum_{n=0}^{\infty} (-1)^n q^{\binom{n+1}{2}} \zeta^{-n} / (1 - \zeta)}{(q; q)_\infty^3} \\ &= \frac{\sum_{n=0}^{\infty} q^{\binom{n+1}{2}} ((-1)^n (\zeta^{-n} - \zeta^{n+1}) / (1 - \zeta))}{\sum_{n=0}^{\infty} (-1)^n (2n + 1) q^{\binom{n+1}{2}}} \\ &\quad \text{(by [9, p. 285, Th. 357])} \\ &= \frac{\sum_{n=0}^{\infty} q^{\binom{n+1}{2}} - 3 \sum_{n=0}^{\infty} q^{\binom{3n+2}{2}}}{\sum_{n=0}^{\infty} (-1)^n (2n + 1) q^{\binom{n+1}{2}}}. \end{aligned}$$

□

From (3.4) we immediately deduce a computationally fast recurrence for $PD(n)$.

Corollary 6. For $n \geq 0$

$$(3.9) \quad \sum_{j \geq 0} PD \left(n - \binom{j+1}{2} \right) (-1)^j (2j+1) = \begin{cases} 0 & \text{if } n \neq \text{triangular number} \\ 1 & \text{if } n = \binom{m+1}{2} \text{ and } m \not\equiv 1 \pmod{3} \\ -2 & \text{otherwise} \end{cases}$$

Corollary 7. $PD(3n + 2) \equiv 0 \pmod{3}$.

Proof: By (3.4), we see that

$$\begin{aligned} pd(q) &\equiv \frac{\sum_{n=0}^{\infty} q^{\binom{n+1}{2}}}{(q; q)_{\infty}^3} \pmod{3} \\ &\equiv \frac{\sum_{n=0}^{\infty} q^{\binom{n+1}{2}}}{(q^3; q^3)_{\infty}} \pmod{3} \end{aligned}$$

but this last expression has no powers of $q \equiv 2 \pmod{3}$ in its expansion. \square

Theorem 8. For $|q| < 1$ and $\zeta = e^{\pi i/3}$,

$$(3.10) \quad pdo(q) = \frac{(\zeta q; q^2)_{\infty} (\zeta^{-1} q; q^2)_{\infty}}{(q; q^2)_{\infty}^2}$$

$$(3.11) \quad = 1 + \sum_{n=1}^{\infty} \frac{(-q^6; q^6)_{n-1} q^n}{(q; q)_{2n} (-q^2; q^2)_{n-1}}$$

$$(3.12) \quad = 1 - \zeta \sum_{n=1}^{\infty} \frac{(\zeta q; q)_{2n-1} (\zeta q^2; q^2)_{n-1} (-1)^n q^{n^2}}{(q; q)_{2n} (q; q^2)_n}$$

$$(3.13) \quad = \frac{\sum_{n=-\infty}^{\infty} q^{(3n)^2} - \sum_{n=-\infty}^{\infty} q^{(3n+1)^2}}{1 + 2 \sum_{n=1}^{\infty} (-1)^n q^{n^2}}.$$

Proof: By Corollary 3 with $k = 2$

$$(3.14) \quad pdo(q) = pd(q)/pd(q^2)$$

$$(3.15) \quad = \frac{(\zeta q; q)_{\infty} (\zeta^{-1} q; q)_{\infty}}{(q; q)_{\infty}^2} \cdot \frac{(q^2; q^2)_{\infty}}{(\zeta q^2; q^2)_{\infty} (\zeta^{-1} q^2; q^2)_{\infty}}$$

$$(3.16) \quad = \frac{(\zeta q; q^2)_{\infty} (\zeta^{-1} q; q^2)_{\infty}}{(q; q^2)_{\infty}^2}.$$

Identity (3.11) may be rewritten as follows:

$$(3.17) \quad \sum_{n=0}^{\infty} \frac{(\zeta; q^2)_n (\zeta^{-1}; q^2)_n}{(q^2; q^2)_n (q; q^2)_n} q^n = \frac{(\zeta^{-1} q; q^2)_{\infty} (\zeta q; q^2)_{\infty}}{(q; q^2)_{\infty} (q; q^2)_{\infty}},$$

and in this form we see that it is a specialization of the q -analog of Gauss's theorem [2, p. 20, Cor. 2.4, q replaced by q^2 , $a = \zeta$, $b = \zeta^{-1}$, $c = q$].

Identity (3.12) may be rewritten as follows:

$$\begin{aligned} \lim_{t \rightarrow 0} \sum_{n=0}^{\infty} \frac{(\zeta/q; q^2)_n (1 - \zeta q^{2n-1}) (\zeta; q^2)_n^2 (t^{-1}; q^2)_n}{(q^2; q^2)_n (1 - \zeta/q) (q; q^2)_n^2 (\zeta q t; q^2)_n} t^n q^n \zeta^{-n} \\ = \frac{(\zeta q; q^2)_{\infty} (\zeta^{-1} q; q^2)_{\infty}}{(q; q^2)_{\infty} (q; q^2)_{\infty}}, \end{aligned}$$

and in this form we see that it is a specialization of the limiting form of Jackson's theorem [6, p. 238, eq. (II.20)].

To obtain (3.13) we note that

$$(3.18) \quad (-\zeta)^n + (-\zeta)^{-n} = \begin{cases} 2 & \text{if } n \equiv 0 \pmod{3} \\ -1 & \text{otherwise.} \end{cases}$$

Hence by Jacobi's triple product

$$\begin{aligned} \frac{(\zeta q; q^2)_{\infty} (\zeta^{-1} q; q^2)_{\infty}}{(q; q^2)_{\infty}^2} &= \frac{(q^2; q^2)_{\infty} (\zeta q; q^2)_{\infty} (\zeta^{-1} q; q^2)_{\infty}}{(q^2; q^2)_{\infty} (q; q^2)_{\infty}^2} \\ &= \frac{\sum_{n=-\infty}^{\infty} (-1)^n q^{n^2} \zeta^n}{\sum_{n=-\infty}^{\infty} (-1)^n q^{n^2}} \\ &= \frac{1 + \sum_{n=1}^{\infty} q^{n^2} ((-\zeta)^n + (-\zeta)^{-n})}{1 + 2 \sum_{n=1}^{\infty} (-1)^n q^{n^2}} \\ &= \frac{\sum_{n=-\infty}^{\infty} q^{(3n)^2} - \sum_{n=-\infty}^{\infty} q^{(3n+1)^2}}{1 + 2 \sum_{n=1}^{\infty} (-1)^n q^{n^2}}. \end{aligned}$$

□

From (3.13) we immediately deduce a computationally fast recurrence for $PDO(n)$.

Corollary 9. For $n \geq 0$

$$(3.19) \quad PDO(n) + 2 \sum_{j \geq 1} PDO(n - j^2)(-1)^j = \begin{cases} 0 & \text{if } n \text{ is not a square} \\ 1 & \text{if } n = 0 \\ 2 & \text{if } n = (3m)^2 \\ -1 & \text{otherwise.} \end{cases}$$

Corollary 10. $PDO(n)$ is odd precisely when either n is 0 or a square not divisible by 3.

Proof: By (3.13)

$$pdo(q) \equiv 1 - \sum_{n=-\infty}^{\infty} q^{(2n+1)^2} \pmod{2}.$$

□

4. RELATIONS TO OTHER PARTITION FUNCTIONS

In this section we shall consider three classical partition functions with an extensive literature.

$M(q)$ will denote the generating function for partitions without 1's and without any two parts differing by 1. It is a theorem of MacMahon [12, p.54] (see also [1]) that

$$(4.1) \quad M(q) = \prod_{\substack{n \neq \pm 0, 1 \\ (\text{mod } 6)}}^{\infty} (1 - q^n)^{-1} = \frac{(-q^3; q^3)_{\infty}}{(q^2; q^2)_{\infty}}$$

$S(q)$ will denote the generating function for partitions in which parts differ by at least 3 and multiples of 3 differ by at least 4. Schur [16] has shown that

$$(4.2) \quad S(q) = \prod_{\substack{n \neq \pm 2, 3 \\ (\text{mod } 6)}}^{\infty} (1 - q^n)^{-1} = \frac{(-q; q)_{\infty}}{(-q^3; q^3)_{\infty}}.$$

Our final generating function, $\Phi_2(q)$, is related to more complicated objects, generalized Frobenius partitions, and we refer the reader to [3] for an introduction of this topic. $\Phi_2(q)$ is the generating function for generalized Frobenius partitions wherein any number can appear on each row as a part

at most twice. In [3, eq. (5.11)], it is shown that

$$\begin{aligned} \Phi_2(q) &= \prod_{n=1}^{\infty} \frac{1}{(1-q^n)(1-q^{12n-10})(1-q^{12n-2})(1-q^{12n-3})(1-q^{12n-2})} \\ &= \frac{\sum_{m=-\infty}^{\infty} q^{(3m)^2} - \sum_{m=-\infty}^{\infty} q^{(3m+1)^2}}{(q; q)_{\infty}^2} \\ &= \frac{1}{(q; q)_{\infty}^2} \quad (\text{by [3, eq. (10.4)]}). \end{aligned}$$

Given the above generating functions, we can easily combine these formulas with those of Sections 2 and 3 to deduce the following:

Theorem 11. For $|q| < 1$,

$$(4.3) \quad pd(q) (q; q)_{\infty} = M(q)$$

$$(4.4) \quad pd(q) S(q) = \frac{1}{(q; q)_{\infty}^2}$$

$$(4.5) \quad pdo(q) = (q^2; q^2)_{\infty} \Phi_2(q)$$

Proof: Identity (4.3) follows from (4.1) and (1.3); identity (4.4) follows from (4.2) and (1.3). Finally identity (4.5) follows from (4.3) and (3.13). \square

It should be observed that these relationships are quite simple and suggest the possibility of our obtaining insight for partitions with designated summands from these classical partition functions.

5. ARITHMETIC PROPERTIES

Because of their relationships with eta functions and modular forms, partition functions typically feature many interesting arithmetic properties. Our work on partitions with designated summands has produced infinite families of partitions whose generating functions are products of eta functions, and in this section we will give a sample of some divisibility and congruence properties for some of these partition functions. To work in the context of modular forms, we require the following facts:

Proposition 12 (Gordon, Hughes, Newman, Ligozat). *Let*

$$f(z) = \prod_{1 \leq \delta | N} \eta^{r_{\delta}}(\delta z)$$

be a product of eta functions which satisfies the following criteria

(i)

$$\sum_{\delta | N} \delta r_{\delta} \equiv 0 \pmod{24}$$

(ii)

$$\sum_{\delta|N} \frac{N}{\delta} r_\delta \equiv 0 \pmod{24}$$

(iii) For each $d|N$,

$$\sum_{\delta|N} (d, \delta)^2 \frac{r_\delta}{\delta} \geq 0$$

Then, $f(z) \in M_k(\Gamma_0(N), \chi)$ if $k = \frac{1}{2} \sum r_\delta \in \mathbb{N}$ and χ is the quadratic character defined by

$$(5.1) \quad \chi(\ell) = \left(\frac{(-1)^k \prod \delta^{r_\delta}}{\ell} \right)$$

Here $M_k(\Gamma_0(N), \chi)$ is the \mathbb{C} - vector space of holomorphic modular forms of weight k , level N , and Nebentypus character χ . Conditions (i)-(ii) ensure that $f\left(\frac{az+b}{cz+d}\right) = \chi(d)(cz+d)^k f(z)$ for all z with $\text{Im}(z) > 0$ and for all integers a, b, c, d such that $ad - bc = 1$ and $N|c$. Condition (iii) guarantees that f is "holomorphic at the cusps." For more on the basic theory of modular forms, see Koblitz [10]. We recall the following well-known facts about modular forms and their Fourier expansions.

Proposition 13. [10] Suppose that $f(z) \in M_k(\Gamma_0(N), \chi)$ with Fourier expansion $f(z) = \sum_{n=0}^{\infty} a(n)q^n$. For any positive integer t ,

(i)

$$f(z) | T(t) := \sum_{n=0}^{\infty} \sum_{d|\gcd(t,n)} \chi(d) d^{k-1} a\left(\frac{tn}{d^2}\right) q^n$$

is the Fourier expansion of a modular form in $M_k(\Gamma_0(N), \chi)$.

(ii)

$$f(z) | V(t) := \sum_{n=0}^{\infty} a(n) q^{tn}$$

is the Fourier expansion of a modular form in $M_k(\Gamma_0(tN), \chi)$.

The operators $T(t)$ are the Hecke operators for integral weight modular forms. Note that as a special case of part (i) of this proposition we have that for any $t|N$,

$$f(z) | U(t) := \sum_{n=0}^{\infty} a(tn) q^n$$

is the Fourier expansion of a modular form in $M_k(\Gamma_0(N), \chi)$.

Proposition 14. [18] Suppose $f(z) = \sum_{n=0}^{\infty} a(n)q^n \in M_k(\Gamma_0(N), \chi)$ satisfies

(i) $a(n) \in \mathbb{Z}$ for all n

(ii) $a(n) \equiv 0 \pmod{M}$ for all $n \leq 1 + \frac{kN}{12} \prod_{p|N} (1 + \frac{1}{p})$

Then, $a(n) \equiv 0 \pmod{M}$ for all n .

Lemma 15. Suppose that $f = \sum_{n=0}^{\infty} a(n)q^n$, $g = \sum_{n=0}^{\infty} d(mn)q^{mn}$ with $d(0) = 1$, and

$$fg = \sum_{n=0}^{\infty} b(n)q^n$$

Then,

- (i) If $a(mn + r) \equiv 0 \pmod{M}$ for $0 \leq n \leq B$, then $b(mn + r) \equiv 0 \pmod{M}$ for $0 \leq n \leq B$.
- (ii) If $b(mn + r) \equiv 0 \pmod{M}$ for all n , then $a(mn + r) \equiv 0 \pmod{M}$ for all n .

Proof: Both facts follow quickly after writing the coefficient of q^{mn+r} from fg in two ways:

$$b(mn + r) = a(mn + r) + \sum_{k \geq 1} d(mk)a(m(n - k) + r)$$

Part (i) is obvious and part (ii) comes from a simple induction argument. \square

We begin with a theorem about the divisibility of $PDp(n)$ when p is prime. The notation $PDp(n)$ is for the number of partitions of n with designated summands wherein no part is divisible by p . Recall that we say *almost all* natural numbers have a property P if

$$\lim_{N \rightarrow \infty} \frac{\#\{n \leq N : P(n)\}}{N} = 1.$$

Theorem 16. If p is any prime and r any nonnegative integer, then almost all n have the property that $PDp(n)$ is divisible by p^r

Proof: It is a theorem of Serre [17] that for any positive integer m , almost all of the Fourier coefficients of an integer weight holomorphic modular form are divisible by m . Hence the theorem will follow if we can find, for each p and sufficiently large r , modular forms $F_{r,p}(z)$ such that

$$F_{r,p}(z) \equiv \sum_{n=0}^{\infty} PDp(n)q^n \pmod{p^r}$$

To this end we define for all r, p with $r \geq 1$ and $(p, 6) = 1$,

$$F_{r,p}(z) := \frac{\eta(6z)\eta(pz)\eta(2pz)\eta(3pz)\eta^{p^r}(z)}{\eta(z)\eta(2z)\eta(3z)\eta(6pz)\eta^{p^{r-1}}(pz)}$$

One can demonstrate that

$$F_{r,p}(z) \in M_{\frac{p^{r-1}(p-1)}{2}}(\Gamma_0(72p), \chi)$$

(where the character χ is defined in Proposition 12) by verifying the conditions of Proposition 12: It is easy to find that

$$\sum_{\delta|72p} \delta r_\delta = 0$$

and

$$\sum_{\delta|72p} \frac{72p}{\delta} r_\delta = 24(5 - 5p + 3p^{r+1} - 3p^{r-1})$$

For condition (iii), notice that for any $d|72p$,

$$\sum \frac{(d, \delta)^2 r_\delta}{\delta}$$

depends only on $(d, 6p)$. In the table below are shown the divisors of $6p$ and the corresponding sums. They are all nonnegative which completes the proof for $p \neq 2, 3$.

$d 6p$	$\sum \frac{(d, \delta)^2 r_\delta}{\delta}$
$d = 1$	$\frac{5(1-p)}{3p} + p^r - p^{r-2}$
$d = 2$	$\frac{8(1-p)}{3p} + p^r - p^{r-2}$
$d = 3$	$\frac{3(1-p)}{p} + p^r - p^{r-2}$
$d = 6$	$p^r - p^{r-2}$
$d = p$	$\frac{5(p-1)}{3}$
$d = 2p$	$\frac{8(p-1)}{3}$
$d = 3p$	$3(p-1)$
$d = 6p$	0

We shall omit the details for the cases $p = 2, 3$ which do not quite fit the argument above. Still it is routine to show that

$$\begin{aligned} G_{r,2}(z) &:= \frac{\eta^2(6z)\eta(4z)\eta^{2^r}(z)}{\eta(z)\eta(3z)\eta(12z)\eta^{2^{r-1}}(2z)} \\ &\equiv \sum_{n=0}^{\infty} PDO(n)q^n \pmod{2^r} \end{aligned}$$

is in $M_{2^{r-2}}(\Gamma_0(144), \chi_{triv})$ for $r \geq 2$ and that

$$\begin{aligned} H_{r,3}(z) &:= \frac{\eta^2(6z)\eta(9z)\eta^{3^r}(z)}{\eta(z)\eta(2z)\eta(18z)\eta^{3^{r-1}}(3z)} \\ &\equiv \sum_{n=0}^{\infty} PD3(n)q^n \pmod{3^r} \end{aligned}$$

is in $M_{3^{r-1}}(\Gamma_0(108), \chi)$ for all r . \square

Although this theorem already contains a strong statement about $PDO(n)$ modulo powers of 2, one can obtain more specific information using Hecke operators. For example, we find by an application of Sturm's criterion that

$$pdo(q) \mid T(p) \equiv 0 \pmod{4}$$

for many small primes including $p = 7, 11, 19, 23, 31$. This implies that

$$PDO(pn) - PDO(n/p) \equiv 0 \pmod{4}$$

for all such primes p and all nonnegative integers n , which immediately gives infinitely many congruences in arithmetic progressions modulo 4. The same techniques can be employed to produce specific information about any of the functions $pdp(q)$ modulo p^r , if desired.

Next we focus on some congruence properties of partitions with designated summands. We shall demonstrate a technique that can be used to computationally verify congruences for these partition functions and give some examples for the generating functions $pd(q)$ and $pdo(q)$. The first step is to combine Propositions 13 and 14 and Lemma 15 into a useful criterion. This technique was employed by Eichhorn and Ono [4] to prove congruence properties of modular functions.

Theorem 17. *Suppose that*

$$\begin{aligned} f(z) &:= \sum_{n=0}^{\infty} a(n)q^n \times \prod \eta^{r_{t\delta}}(t\delta z) \\ &:= \sum_{n=0}^{\infty} r(n)q^n \\ &\in M_k(\Gamma_0(N), \chi) \end{aligned}$$

and that $a(tn - b) \equiv 0 \pmod{M}$ for all

$$n \leq B := 1 + \frac{kN}{12} \prod_{p|N} \left(1 + \frac{1}{p}\right)$$

where $b = \frac{\sum t\delta r_{t\delta}}{24}$. Then, $a(tn - b) \equiv 0 \pmod{M}$ for all n .

Proof: By the remark following Proposition 13,

$$f(z) | U(t) = \sum_{n=0}^{\infty} r(tn)q^n \in M_k(\Gamma_0(N), \chi)$$

so $r(tn) \equiv 0 \pmod{M}$ for all n if $r(tn) \equiv 0 \pmod{M}$ for all $n \leq B$ by Proposition 14. Now, by the first part of Lemma 15, this will happen if $a(tn - b) \equiv 0 \pmod{M}$ for all $n \leq B$ which in turn will imply that $a(tn - b) \equiv 0 \pmod{M}$ for all n by the second part of Lemma 15. \square

The above theorem successfully transfers Sturm's criterion from modular forms to other series under suitable conditions. We begin with one detailed argument.

Corollary 18. *For all nonnegative integers n ,*

$$(5.2) \quad PD(12n + 10) \equiv 0 \pmod{8}$$

Proof: Define the η - product $f(z)$ by

$$\begin{aligned} f(z) &:= \frac{\eta(6z)\eta^{16}(12z)\eta^3(24z)\eta^4(36z)\eta^7(72z)}{\eta(z)\eta(2z)\eta(3z)} \\ &= \sum_{n=0}^{\infty} PD(n)q^n \times \eta^{16}(12z)\eta^3(24z)\eta^4(36z)\eta^7(72z) \\ &\in M_{14}(\Gamma_0(72), \chi_f) \end{aligned}$$

according to Proposition 12. In the notation of theorem 17,

$$b = \frac{12 \cdot 16 + 3 \cdot 24 + 4 \cdot 36 + 7 \cdot 72}{24} = 38$$

and hence if $PD(12n - 38) \equiv 0 \pmod{8}$ for all $n \leq 169$ then $PD(12n - 38) \equiv 0 \pmod{8}$ for all n . It has been verified computationally that $PD(12n + 10) \equiv 0 \pmod{8}$ for all $n \leq 165$ which implies the congruence (5.2) for all n . \square

We have found numerous congruences for partitions with designated summands which can be proven computationally using exactly the same argument as in Corollary 18. The work is in finding the right modular form $f(z)$ which implies the desired congruence by an application of Theorem 17. Below we list many congruence properties and the weight k , level N , holomorphic modular forms $f(z)$ used to verify the congruences.

Corollary 19. *The following congruences are implied by the existence of the corresponding holomorphic modular forms $f(z)$ and a sufficient computation.*

Congruence	$f(z) \in M_k(\Gamma_0(N), \chi_f)$	(k, N)
$PD(24n + 18) \equiv 0 \pmod{24}$	$\frac{\eta(6z)\eta^{27}(24z)\eta^{24}(48z)\eta(72z)\eta^4(144z)}{\eta(z)\eta(2z)\eta(3z)}$	(27, 144)
$PD(12n + 6) \equiv 0 \pmod{4}$	$\frac{\eta(6z)\eta^{16}(12z)\eta^4(24z)\eta^4(36z)\eta^4(72z)}{\eta(z)\eta(2z)\eta(3z)}$	(13, 72)
$PDO(9n + 6) \equiv 0 \pmod{3}$	$\frac{\eta^2(6z)\eta(4z)\eta^8(9z)\eta^8(54z)\eta^2(108z)}{\eta(z)\eta(3z)\eta(12z)}$	(9, 108)
$PDO(12n + 10) \equiv 0 \pmod{3}$	$\frac{\eta^2(6z)\eta(4z)\eta^{12}(12z)\eta^4(24z)\eta^8(48z)}{\eta(z)\eta(3z)\eta(12z)}$	(12, 48)
$PDO(12n + 6) \equiv 0 \pmod{3}$	$\frac{\eta^2(6z)\eta(4z)\eta^{12}(12z)\eta^{12}(72z)}{\eta(z)\eta(3z)\eta(12z)}$	(12, 72)
$PDO(24n + 16) \equiv 0 \pmod{3}$	$\frac{\eta^2(6z)\eta(4z)\eta^{16}(24z)\eta^8(48z)}{\eta(z)\eta(3z)\eta(12z)}$	(12, 48)
$PDO(24n) \equiv 0 \pmod{3}$	$\frac{\eta^2(6z)\eta(4z)\eta^{24}(24z)}{\eta(z)\eta(3z)\eta(12z)}$	(12, 144)

There are undoubtedly many more simple congruences for the functions listed above and indeed for every member of our infinite families of generating functions which are η - products. The profusion of these arithmetic properties, especially for small moduli, suggests the likelihood of finding combinatorial explanations. Such a revelation could perhaps lead to the discovery and classification of families of congruences and a deeper understanding of partitions with designated summands.

6. MODULAR RELATIONS

In this section we consider the generating functions for $PD(n)$ and $PDO(n)$ in certain arithmetic progressions. We show that many of these generating functions are infinite products and also demonstrate that the phenomenon in Andrews' tenth problem [3, 5] on Frobenius partitions arises in the study of partitions with designated summands. Each of the Theorems 21 – 24 was observed by applying Euler's algorithm for infinite products [2, p. 98]. An application of Sturm's criterion guarantees that our observation was, in fact, a proof.

Theorem 20.

$$\sum_{n=0}^{\infty} PD(2n)q^n = \frac{(q^4; q^4)_{\infty}^2 (q^6; q^6)_{\infty}^5}{(q; q)_{\infty}^3 (q^2; q^2)_{\infty} (q^3; q^3)_{\infty}^3 (q^{12}; q^{12})_{\infty}^2}$$

$$\sum_{n=0}^{\infty} PD(2n+1)q^n = \frac{(q^2; q^2)_{\infty}^5 (q^{12}; q^{12})_{\infty}^2}{(q; q)_{\infty}^5 (q^3; q^3)_{\infty} (q^4; q^4)_{\infty}^2 (q^6; q^6)_{\infty}}$$

Proof: By Propostion 12, the equation

$$\frac{\eta^{11}(2z)\eta(6z)}{\eta(z)\eta(3z)} = \frac{\eta^9(2z)\eta^2(8z)\eta^5(12z)}{\eta(4z)\eta^3(6z)\eta^2(24z)} + \frac{\eta^7(2z)\eta^5(4z)\eta^2(24z)}{\eta(6z)\eta^2(8z)\eta(12z)}$$

is an equation between modular forms in $M_5(\Gamma_0(72), (\frac{-1}{\cdot}))$ and hence is verified by comparing the first 61 terms on each side. Multiply both sides by $\eta^{-12}(2z)$ and we have

$$\begin{aligned} \sum_{n=0}^{\infty} PD(2n)q^{2n} + \sum_{n=0}^{\infty} PD(2n+1)q^{2n+1} &= \frac{\eta(6z)}{\eta(z)\eta(2z)\eta(3z)} \\ &= \frac{(q^8; q^8)_{\infty}^2 (q^{12}; q^{12})_{\infty}^5}{(q^2; q^2)_{\infty}^3 (q^4; q^4)_{\infty} (q^6; q^6)_{\infty}^3 (q^{24}; q^{24})_{\infty}^2} \\ &\quad + q \times \frac{(q^4; q^4)_{\infty}^5 (q^{24}; q^{24})_{\infty}^2}{(q^2; q^2)_{\infty}^5 (q^6; q^6)_{\infty} (q^8; q^8)_{\infty}^2 (q^{12}; q^{12})_{\infty}} \end{aligned}$$

and the theorem follows. \square

In an identical fashion we obtain Theorems 21 and 22.

Theorem 21.

$$\begin{aligned} \sum_{n=0}^{\infty} PDO(2n)q^n &= \frac{(q^4; q^4)_{\infty}^2 (q^6; q^6)_{\infty}^4}{(q; q)_{\infty}^2 (q^3; q^3)_{\infty}^2 (q^{12}; q^{12})_{\infty}^2} \\ \sum_{n=0}^{\infty} PDO(2n+1)q^n &= \frac{(q^2; q^2)_{\infty}^6 (q^{12}; q^{12})_{\infty}^2}{(q; q)_{\infty}^4 (q^4; q^4)_{\infty}^2 (q^6; q^6)_{\infty}^2} \end{aligned}$$

Proof: The equation

$$\frac{\eta^{12}(2z)\eta(4z)\eta^2(6z)}{\eta(z)\eta(3z)\eta(12z)} = \frac{\eta^{10}(2z)\eta^2(8z)\eta^4(12z)}{\eta^2(6z)\eta^2(24z)} + \frac{\eta^8(2z)\eta^6(4z)\eta^2(24z)}{\eta^2(8z)\eta^2(12z)}$$

is an equation between modular forms in $M_6(\Gamma_0(144), \chi_{triv})$ and hence is verified by checking the first 145 terms on each side. The claim follows immediately as in Theorem 20 after multiplying both sides by $\eta^{-12}(2z)$.

Theorem 22.

$$\begin{aligned} \sum_{n=0}^{\infty} PDO(3n)q^n &= \frac{(q^2; q^2)_{\infty}^2 (q^6; q^6)_{\infty}^4}{(q; q)_{\infty}^4 (q^{12}; q^{12})_{\infty}^2} \\ \sum_{n=0}^{\infty} PDO(3n+1)q^n &= \frac{(q^2; q^2)_{\infty}^4 (q^3; q^3)_{\infty}^3 (q^{12}; q^{12})_{\infty}}{(q; q)_{\infty}^5 (q^4; q^4)_{\infty} (q^6; q^6)_{\infty}^2} \\ \sum_{n=0}^{\infty} PDO(3n+2)q^n &= 2 \times \frac{(q^2; q^2)_{\infty}^3 (q^6; q^6)_{\infty} (q^{12}; q^{12})_{\infty}}{(q; q)_{\infty}^4 (q^4; q^4)_{\infty}} \end{aligned}$$

Proof: The equation

$$\begin{aligned} \frac{\eta^7(3z)\eta(4z)\eta^2(6z)}{\eta(z)\eta(12z)} &= \frac{\eta^4(3z)\eta^2(6z)\eta^4(18z)}{\eta^2(36z)} + \frac{\eta^3(3z)\eta^4(6z)\eta^3(9z)\eta(36z)}{\eta(12z)\eta^2(18z)} \\ &+ 2 \times \frac{\eta^4(3z)\eta^3(6z)\eta(18z)\eta(36z)}{\eta(12z)} \end{aligned}$$

is easily seen by Proposition 12 to be an equation between modular forms in

$M_4(\Gamma_0(144), \chi_{triv})$ and hence is verified by comparing the first 97 terms on each side. Multiply both sides by $\eta^{-8}(3z)$ and the theorem follows. \square

The generating functions for $PD(3n+0, 1, 2)$ do not have the same nice properties as in the previous theorems. However, in one case we find something striking.

Theorem 23. *Define numbers $a(n)$ uniquely by*

$$\sum_{n=0}^{\infty} PD(3n)q^n = \prod_{n=1}^{\infty} (1 - q^n)^{-a(n)}$$

Then for all positive n we have

$$\begin{aligned} a(6n+1) &= 5 \\ a(6n+3) &= 2 \\ a(6n+5) &= 5. \end{aligned}$$

We remark that the sequence $a(2n)$ begins

$$13, -14, 80, -338, 1741, -8902, 16669, \dots$$

Proof of Theorem.: Our object is to show that

$$\sum_{n=0}^{\infty} PD(3n)q^n = (q; q^6)_{\infty}^{-5} (q^3; q^6)_{\infty}^{-2} (q^5; q^6)_{\infty}^{-5} \times F(q^2)$$

for some series F . By Theorem 5, we know that the η -product

$$f(z) := \frac{\eta^{27}(6z)\eta^4(3z)}{\eta(z)\eta(2z)\eta^3(9z)}$$

is a modular form in $M_{13}(\Gamma_0(18), \chi_f)$, and so is $g(z) := f(z) | U(3)$. Moreover,

$$h(z) := g(z) - g(z) | U(2) | V(2)$$

is a modular form in $M_{13}(\Gamma_0(36), \chi_f)$. An application of Sturm's criterion shows that $h(z) = 0$. In other words, $g(z) = \sum_{n \geq 1} c(n)q^{2n}$. To complete

the proof, observe that

$$\begin{aligned} g(z) &= \frac{\eta^{27}(6z)\eta^4(3z)}{\eta(z)\eta(2z)\eta^3(9z)} \mid U(3) \\ &= pd(q) \mid U(3) \times q^2(q^2; q^2)_\infty^{26} \times \frac{(q; q)_\infty^5}{(q^3; q^3)_\infty^3} \end{aligned}$$

and therefore

$$\begin{aligned} \sum_{n=0}^{\infty} PD(3n)q^n &= pd(q) \mid U(3) \\ &= \frac{(q^3; q^3)_\infty^3}{(q; q)_\infty^5} \times \frac{g(z)}{q^2(q^2; q^2)_\infty^{26}} \\ &= (q; q^6)_\infty^{-5} (q^3; q^6)_\infty^{-2} (q^5; q^6)_\infty^{-5} \times F(q^2) \end{aligned}$$

□

7. CONCLUSION

It should be clear that partitions with designated summands are objects with a great deal of structure. It has not been our purpose to give an exhaustive account, but to offer a sample of their combinatorial, functional, and arithmetic properties. By using various tagging schemes, such as colored tags and restrictions on which parts can be designated, one can obtain many more families of infinite product generating functions. These undoubtedly have as many interesting properties as those discussed here, and in specific cases they have further relationships with generalized Frobenius partitions as well as with other types of partitions with designated summands. Since all of these have generating functions which are essentially eta functions, it should be worthwhile to consider the connections between partitions with designated summands and other objects which are counted by Fourier coefficients of modular forms.

REFERENCES

- [1] G. E. Andrews, *A generalization of a partition theorem of MacMahon*, J. Comb. Th. 3 (1967), 100–101.
- [2] G.E. Andrews, *The Theory of Partitions*, Encycl. Math. and Its Applications, Vol. 2, Addison-Wesley, Reading, 1976 [Reissued: Cambridge University Press, Cambridge, 1985, paperback, 1998].
- [3] G.E. Andrews, *Generalized Frobenius partitions*, Mem. Amer. Math. Soc. 49 1984 No. 301, iv + 44 pp.
- [4] D. Eichhorn and K. Ono, *Congruences for partition functions*, in: Analytic Number Theory, Vol. 1, Progr. Math. 138, Birkhauser, 1996, 309-321.
- [5] Frank Garvan, *Modular Functions, Maple and Andrews' 10th. Problem*, in: Topics in Number Theory (Ahlgren, Andrews and Ono eds.), Kluwer, 1999, 163-179.

- [6] G. Gasper and M. Rahman, *Basic Hypergeometric Series*, Encycl. Math and Its Applications, Vol. 35, Cambridge University Press, Cambridge, 1990.
- [7] J. W. L. Glaisher, *Expressions for the first five powers of the series in which the coefficients are the sums of the divisors of the exponents*, Messenger of Math., No. 171, July 1885.
- [8] B. Gordon and K. Hughes, *Multiplicative properties of η -products II*, Cont. Math. 143 (1993), 415-430.
- [9] G. H. Hardy and E. M. Wright, *An Introduction to the Theory of Numbers* 4th ed., Oxford Univ., Press, London and New York, 1960.
- [10] N. Koblitz, *Introduction to Elliptic Curves and Modular Forms*, Springer-Verlag, Berlin/New York, 1984.
- [11] G. Ligozat, *Courbes modulaires de genre 1*, Vol.43, Bulletin de la Soc. Math. de France, 1975.
- [12] P. A. MacMahon, *Combinatory Analysis*, Vol. 2, Cambridge University Press, London, 1916 [Reprinted: Chelsea, New York, 1960].
- [13] P. A. MacMahon, *Divisors of numbers and their continuations in the theory of partitions*, Proc. London Math. Soc., Ser. 2, 19 (1919), 75–113.
- [14] M. Newman, *Construction and application of a certain class of modular functions II*, Proc. London Math. Soc. (3) 9 (1959), 353-387.
- [15] S. Ramanujan, *On certain arithmetical functions*, Proc. Cambridge Phil. Soc., 22 (1916), 159–184. [Reprinted: Collected Papers of S. Ramanujan, Cambridge Univ. Press, Cambridge, 1927, pp. 136–162 Reissued: Chelsea, New York, 1962]
- [16] I. Schur, *Zur additiven Zahlentheorie*, S.-B. Preuss. Akad. Wiss. Phys.-Math. Kl., 1926, pp. 488–495 [Reprinted: I. Schur, Gessammelte Abhandlungen, Vol. 3, pp. 43–50, Springer, Berlin, 1973].
- [17] J.P. Serre, *Divisibilité de certaines fonctions arithmétiques*, L'Ensign. Math 22 (1976), 227-260.
- [18] J. Sturm, *On the Congruence Properties of Modular Forms*, Springer Lect. Notes in Math. Vol.1240, pp. 275-280, Springer-Verlag, Berlin/New York, 1984.

DEPT. OF MATHEMATICS, PENNSYLVANIA STATE UNIVERSITY, UNIVERSITY PARK,
PA 16802 U.S.A.

E-mail address: `andrews@math.psu.edu`

SCHOOL OF MATHEMATICAL SCIENCES, THE UNIVERSITY OF SUSSEX, BRIGHTON
BN1 9QH, U.K.

E-mail address: `R.P.Lewis@sussex.ac.uk`

DEPT. OF MATHEMATICS, UNIVERSITY OF WISCONSIN, MADISON, WI 53706
U.S.A.

E-mail address: `lovejoy@math.wisc.edu`